

Table 12C
2002 Pediatric Nutrition Surveillance ⁽¹⁾
California
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 8/21/2003

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)					
	Number <2500 g	% >4000 g	%	Number <5th	% <5th	% ≥95th	%	Number 85th-<95th	% ≥95th	%	Number 	%
2002	157,676	6.7	9.1	850,756	4.9	5.3	16.2	334,318	16.3	17.3	483,082	13.1
2001	131,493	6.6	9.1	781,922	5.0	5.3	15.9	306,084	16.3	16.5	445,793	13.7
2000	161,684	6.9	9.4	890,600	5.1	5.3	16.1	363,965	16.2	16.7	550,767	13.3
1999	162,300	6.9	9.6	897,904	5.0	5.2	15.6	375,498	16.3	15.9	582,213	13.0
1998	168,177	6.8	9.4	959,431	4.9	5.2	15.4	398,222	16.4	15.5	622,136	13.0
1997	181,986	7.0	9.2	1,013,964	4.9	5.5	14.7	418,198	15.6	14.8	652,081	13.5
1996	201,276	7.2	9.3	1,005,581	5.0	5.6	14.4	403,710	15.6	14.6	682,102	14.0
1995	204,802	8.9	9.2	1,003,204	5.1	5.5	14.6	386,760	15.5	14.4	664,602	14.6
1994	211,518	9.6	8.9	985,233	5.1	5.6	14.3	368,356	15.5	14.2	647,964	14.6
1993	212,705	9.4	9.0	922,675	5.3	6.0	13.9	321,817	15.2	13.8	583,324	14.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2002 Pediatric Nutrition Surveillance ⁽¹⁾
California
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 8/21/2003

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low		High	Short Stature (5)		Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
1992	210,953	9.7	9.2	841,093	5.4	6.2	13.5	266,554	15.1	13.5	511,283	14.4
1991	200,381	9.9	9.5	762,416	5.5	6.2	13.4	230,018	15.2	13.3	459,127	15.5
1990	179,201	8.5	9.5	551,428	6.1	6.6	13.4	151,281	14.9	13.1	373,394	16.4
1989	136,729	6.8	9.5	531,232	7.1	5.8	14.1	146,057	15.9	13.6	237,143	16.7
1988	677	10.5	8.6	4,111	8.1	7.7	12.9	1,010	16.1	13.8	1,212	13.7
1987	0	*	*	1	*	*	*	0	*	*	3	*
1986	1	*	*	0	*	*	*	0	*	*	0	*
1985	0	*	*	0	*	*	*	0	*	*	1	*
1984	1	*	*	0	*	*	*	0	*	*	0	*
1983	2	*	*	0	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2002 Pediatric Nutrition Surveillance ⁽¹⁾
California
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 8/21/2003

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight						Anemia Low Hb/Hct		
	Low		High	Short Stature (5)		Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			(7)	
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
1982	0	*	*	1,966	6.7	6.9	6.0	716	14.7	8.1	893	9.6
1981	0	*	*	1,672	6.8	7.3	5.7	623	13.2	5.8	792	9.3
1980	0	*	*	10,863	7.2	9.2	7.8	4,341	13.7	9.2	5,068	9.7
1979	0	*	*	6	*	*	*	4	*	*	2	*
1978	1	*	*	0	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2002 Pediatric Nutrition Surveillance (1)
California

Page: 1
Run Date: 10/6/2003

Summary of Trends in Growth and Anemia Indicators
Children Aged 5 to <20 Years (2)
Standard Year (School Age Coding)

Summary of Trends in Growth and Anemia Indicators (3)							
Year	Height and Weight					Anemia Low Hb/Hct	
	Short Stature (4)		Under- weight (5)	Over- weight (5)		(6)	
	Number	% <5th	% <5th	% 85th-<95th	% ≥95th	Number	%
2002	512,143	6.7	2.7	18.1	20.8	440,239	12.0
2001	474,493	6.8	2.7	17.8	19.9	406,633	12.4
2000	574,820	6.9	2.8	17.7	19.7	486,065	12.6
1999	600,105	6.8	2.8	17.5	18.9	512,607	11.9
1998	545,455	6.9	2.8	17.2	17.6	473,018	12.1
1997	507,049	7.0	3.2	16.5	16.4	443,223	12.9
1996	434,626	7.5	3.2	16.3	15.6	414,942	13.2
1995	403,902	8.1	3.3	16.2	15.1	384,195	13.8
1994	368,340	8.5	3.4	15.8	14.4	355,349	13.8
1993	343,122	8.9	3.6	15.6	13.7	336,328	13.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C
2002 Pediatric Nutrition Surveillance (1)
California

Page: 2
Run Date: 10/6/2003

Summary of Trends in Growth and Anemia Indicators
Children Aged 5 to <20 Years (2)
Standard Year (School Age Coding)

Summary of Trends in Growth and Anemia Indicators (3)							
Year	Height and Weight					Anemia Low Hb/Hct	
	Short Stature (4)	Under- weight (5)	Over- weight (5)			(6)	
	Number	% <5th	% <5th	% 85th-<95th	% ≥95th	Number	%
1992	288,597	9.5	3.8	15.2	12.9	289,769	13.1
1991	261,071	9.7	3.7	15.0	12.2	267,621	13.9
1990	177,303	10.5	4.0	14.9	12.2	227,213	15.1
1989	156,879	10.6	3.5	15.1	12.7	125,544	16.2
1988	1,083	14.9	3.5	13.5	14.1	747	14.9
1987	0	*	*	*	*	0	*
1986	0	*	*	*	*	0	*
1985	0	*	*	*	*	0	*
1984	0	*	*	*	*	0	*
1983	0	*	*	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C
2002 Pediatric Nutrition Surveillance (1)
California

Page: 3
Run Date: 10/6/2003

Summary of Trends in Growth and Anemia Indicators
Children Aged 5 to <20 Years (2)
Standard Year (School Age Coding)

Summary of Trends in Growth and Anemia Indicators (3)							
Year	Height and Weight					Anemia Low Hb/Hct	
	Short Stature (4)	Under- weight (5)	Over- weight (5)			(6)	
	Number	% <5th	% <5th	% 85th-<95th	% ≥95th	Number	%
1982	714	7.6	2.9	14.8	6.9	590	10.2
1981	678	7.1	2.2	13.3	6.5	548	8.2
1980	5,943	14.3	4.9	12.3	7.1	4,649	9.3
1979	4	*	*	*	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.